





Project Content

□ Family program

• To enhance the resilience of drug abusers' family members or concerned significant others.







| | Evaluation Results (Output indicators) | | |
|-----------------------|---|---|--|
| | Description of output indicators (with expected result) | Achieved Result | |
| Output Indicator 1 | Provide outreaching activities for 3,600 (person-times) EM at-risk individuals, 1,440 (person-times) EM with specific occupation and 600 (person-times) community members through Mass program | Individual: 5,864 At-work: 1,514 Mass program: 1,080 | |
| Output Indicator 2 | Total number of 468 (head count) EM drug abusers / rehabilitees received counselling with Motivational Interviewing, with at least 180 (head count) new /reactivated cases, among which 288 (person-times) case plan set for treatment and rehabilitation | New/re-activated cases: 463 Case plans set with participants: 346 | |
| Output Indicator 3 | Total number of 90 (head count) drug abusers referred to community-based treatment, SACs, DTRCs | 90 | |
| Output Indicator 4 | Total number of 2,160 (person-times) drug abusers / rehabilitees attended the Rehabilitation Strengthening Program & Pre-vocational training | 3,775 | |
| | | | |

| | Evaluation Results (Outcon | ne indicators) |
|-----------------------|---|--|
| | Description of output indicators (with expected result) | Achieved Result |
| Output Indicator 1 | Reduction in drug use frequency among drug abusers (50% of drug abusers reduced drug use after completion of treatment) | 61% (Out of 87 persons completed the treatment) |
| Output Indicator 2 | 40% of case plan achieved among drug abusers / rehabilitees | 73% (Out of 346 case plans) |
| Output Indicator 3 | 50% of participants show improvement in self- efficacy to refuse drug use (or significant improvement in participant's self-efficacy to refuse drug use as indicated by paired t-test) | 60% (Out of 117 respondents) |
| Output Indicator 4 | 40% of participants who attended family training show improvement in capacity to support drug abusers (or significant improvement in participant's capacity to support drug abusers as indicated by paired t-test) | 71% (Out of 152 respondents) |
| | | |





